

LONGEVITY RECORDS OF FLIGHTLESS CORMORANTS

PHALACROCORAX HARRISI

GUSTAVO JIMÉNEZ-UZCÁTEGUI¹, CARLOS A. VALLE^{1,2} & F. HERNÁN VARGAS^{1,3}

¹Department of Sciences, Charles Darwin Foundation, Puerto Ayora, Galápagos, Ecuador (gustavo.jimenez@fcdarwin.org.ec)

²Galápagos Academic Institute for the Arts and Sciences, Universidad San Francisco de Quito, P.O. Box 17-12-841, Quito, Ecuador

³The Peregrine Fund, 5668 West Flying Hawk Lane, Boise, ID 83709, USA

Received 5 March 2012, accepted 18 September 2012

The Flightless Cormorant *Phalacrocorax harrisi* is endemic to the Galápagos Islands, Ecuador (Harris 1973). This species is morphologically and behaviorally unique among extant species of cormorants (Phalacrocoracidae) because of its flightlessness and sequential polyandrous mating system (Valle 1994, Kennedy *et al.* 2009, Valle 2009). The IUCN listed the Flightless Cormorant as endangered (EN) until 2010, but changed its listing in 2011 to vulnerable (VU) “because evidence suggests that the population is stabilizing and the recorded fluctuations during the last three decades have not been extreme” (IUCN 2011). However, its small population size, with numbers fluctuating between 700 and 1 900 individuals during the last three decades (Jiménez-Uzcátegui 2010) as well as its restricted distribution range around Fernandina and the Isabela islands (except the southern and southeastern parts) make the species vulnerable to the impact of recurrent El Niño events (Valle & Coulter 1987, Valle 1995), increases in pathogens (Travis *et al.* 2006, Merkel *et al.* 2007, Deem *et al.* 2010) and introduced predators, such as rats *Rattus* spp. and feral cats *Felis catus*, that prey especially on chicks and juveniles (Valle 1986, 1995).

Of 38 species of cormorants worldwide (Johnsgard 1993), the oldest longevity records for the two species genetically closest to the Flightless Cormorant are 22.6 years for the Double-crested Cormorant *P. auritus*, and 11.9 years for the Neotropic Cormorant *P. brasilianus* (Lutmerding & Love 2011). These two species are considered of least concern (LC) because they have a large

distributional range with large and increasing populations of mature individuals (IUCN 2011).

On 8 April 2006, the monitoring team of one of us (FHV) captured a female Flightless Cormorant *Phalacrocorax harrisi* that was at least 17.2 years old at Zone 4 (0°19'S, 91°23'W) on the northeastern coast of Fernandina Island, Galapagos Archipelago, Ecuador. This cormorant had a metal ring No. 100-029 and had been banded as an adult by one of us (CAV) on 10 July 1990 on South Colony, approximately 3 km southwest of Punta Espinosa, Fernandina Island (Valle 1994); this indicated an actual banded age of 15.9 years. At the time of banding, this bird was sexually mature, therefore at least 17 months old (Valle 1994, 1995, Larrea 2007). This finding establishes the longest life span ever recorded for a Flightless Cormorant. We also provide four additional longevity records for this species based on birds banded by one of us (CAV) between 1989 and 1991 (Table 1).

At present, we are unable to assess whether ages of more than 10 years for Flightless Cormorant are the exception or the norm in their population. As these records were established between 1989 and 2006 (Table 1), we conclude that these individuals are survivors of the last strong El Niño-famine event in 1997–1998. Like many seabird species, Flightless Cormorants have low reproductive levels and depends directly on food availability, which, in turn, depends on oceanographic conditions (Harris 1979, Tindle 1984,

TABLE 1
Longevity records of Flightless Cormorants *Phalacrocorax harrisi*, Galápagos Islands

Identification			Banding					Re-capture				Age (years)	
Pit-tag	Band No.	Sex ^a	Date	Age ^b	Breeding Status	Locality	Coordinates	Date	Breeding Status	Locality	Coordinates	Banded	Estimated
53812301	100029	F	10-Jul-1990	A	Courtship	South Colony	0°17'12.08'S 91°25'5.58'W	08-Apr-2006	Incubation eggs	Zone 4	0°19'45.30'S 91°23'41.64'W	15.9	17.2
53512101	100035	F	26-Jul-1990	A	Nesting chicks	Espinoza	0°15'50.61'S 91°26'40.52'W	02-Oct-2003	Solitary	Serrano	0°17'12.08'S 91°25'5.58'W	13.3	14.8
42023011	100060	F	21-Nov-1991	J	Solitary	Valle	0°15'38.55'S 91°27'34.27'W	08-Apr-2006	Solitary	Valle	0°15'38.55'S 91°27'34.27'W	14.5	14.7
53380633	100043	M	28-Jul-1990	A	Nest building	South Colony	0°17'12.08'S 91°25'5.58'W	26-Aug-2003	Solitary	Serrano	0°17'12.08'S 91°25'5.58'W	13.1	14.6
42045262	100130	M	29-Jun-1989	A	Nesting juvenile	Valle	0°15'38.55'S 91°27'34.27'W	22-Apr-2001	Solitary	Escondida	0°15'45.25'S 91°28'7.21'W	11.1	12.6

^a F: Female, M: Male

^b A: Adult, J: Juvenile

Valle 1994). Hence, the protection of adult cormorants surviving El Niño events should be top priority, as they have the potential to reproduce and restore the diminished populations immediately after such extreme climatic events. These records also reinforce the importance of preventing cumulative increases in adult mortality from disease, predation, entanglement in fishing gear and other anthropogenic threats. Therefore, we recommend maximizing survival of adult cormorants as a protection measure to achieve long-term conservation goals for the species.

ACKNOWLEDGEMENTS

We thank the Galápagos National Park Service and the Charles Darwin Foundation. We wish to thank the donors at that time including, among others, Princeton University, National Geographic, Swarovski & Co., Darwin Initiative, Galápagos Conservancy, SeaWorld & Busch Gardens Conservation Fund and the Swiss Friends of Galapagos Organization (FOGO). We are grateful to P. Piedrahita who captured the oldest cormorant, as well as to C. Larrea, S. Naranjo and F. Gil, collaborators and volunteers who worked in the Cormorant and Penguin Program. We thank C. Nelson Gallardo for grammar correction on an earlier version and for the commentaries from the referees in this manuscript. This is contribution number 2051 of the Charles Darwin Foundation for the Galápagos Islands.

REFERENCES

- DEEM, S.L., MERKEL, J., BALLWEBER, L., VARGAS, F.H., CRUZ, M. & PARKER, P.G. 2010. Exposure to *Toxoplasma gondii* in Galápagos Penguins (*Spheniscus mendiculus*) and Flightless Cormorants (*Phalacrocorax harrisi*) in the Galápagos Islands, Ecuador. *Journal of Wildlife Diseases* 46: 1005–1011.
- HARRIS, M.P. 1973. The Galápagos Avifauna. *Condor* 75: 265–278.
- HARRIS, M.P. 1979. Population dynamics of the Flightless Cormorant *Nannopterum harrisi*. *Ibis* 121: 135–146.
- IUCN. 2011. IUCN Red List of Threatened Species. Version 2010.3. www.iucnredlist.org, accessed 1 June 2011.
- JIMÉNEZ-UZCÁTEGUI, G. 2010. Monitoreo del Pingüino de Galápagos y Cormorán no volador 2010. Puerto Ayora, Ecuador: Informe técnico para la Fundación Charles Darwin y Parque Nacional Galápagos.
- JOHNSGARD, P.A. 1993. Cormorants, darters, and pelicans of the world. Washington: Smithsonian Institution Press.
- KENNEDY, M., VALLE, C.A. & SPENCER, H.G. 2009. The phylogenetic position of the Galápagos Cormorant. *Molecular Phylogenetics and Evolution* 53: 94–98.
- LARREA, C. 2007. Movimiento, dispersión y éxito reproductivo del cormorán no volador *Phalacrocorax harrisi* en las islas Galápagos. [BSc biology thesis]. Pontificia Universidad Católica del Ecuador, Quito, Ecuador.
- LUTMERDING, J.A. & LOVE, A.S. 2011. Longevity records of North American birds. Version 2011.2. Laurel, Maryland: Patuxent Wildlife Research Center, Bird Banding Laboratory. http://www.pwrc.usgs.gov/bbl/longevity/longevity_main.cfm, accessed 24 November 2011.
- MERKEL, J., JONES, H.I., WHITEMAN, N.K., GOTTDENKER, N., VARGAS, F.H., TRAVIS, E.K., MILLER, R.E. & PARKER, P.G. 2007. Microfilariae in Galápagos Penguins (*Spheniscus mendiculus*) and Flightless Cormorants (*Phalacrocorax harrisi*): Genetics, morphology and prevalence. *Journal of Parasitology* 93: 495–503.
- TINDLE, R.W. 1984. The evolution of breeding strategies in the Flightless Cormorant (*Nannopterum harrisi*) of the Galapagos Islands. *Proceedings of the Linnean Society of London* 21: 157–164.
- TRAVIS, E.K., VARGAS, F.H., MERKEL, J., GOTTDENKER, N., MILLER, R.E. & PARKER, P.G. 2006. Hematology, serum chemistry and serology of the Flightless Cormorant (*Phalacrocorax harrisi*) in the Galápagos Islands, Ecuador. *Journal of Wildlife Diseases* 42: 133–141.
- VALLE, C.A. 2009. The Flightless Cormorant: the evolution of female rule. In: De Roy, T. (Ed.). Galápagos: preserving Darwin's legacy. New Zealand: D. Bateman, Ltd. pp. 162–169.
- VALLE, C.A. 1995. Effective population size and demography of the rare Flightless Galápagos Cormorant. *Ecological Applications* 5: 601–617.
- VALLE, C.A. 1994. The ecology and evolution of sequential polyandry in Galápagos Cormorants (*Compsahallieus [Nannopterum] harrisi*). [PhD thesis]. Princeton University, Princeton, NJ.
- VALLE, C.A. 1986. Status of the Galápagos Penguin and Flightless Cormorant. *Noticias de Galápagos* 43: 16–17.
- VALLE, C.A. & Coulter, M.C. 1987. Present status of the Flightless Cormorant, Galápagos Penguin and Greater Flamingo populations in the Galápagos Islands, Ecuador, after the 1982–1983 El Niño. *Condor* 89: 276–281.