FROM COAST TO DESERT: FIRST INLAND RECORD OF THE SOOTY GULL *ICHTHYAETUS HEMPRICHII* IN SAUDI ARABIA

ABDELWAHAB CHEDAD[1]*, ABDULAZIZ MUBARAK AL-QAHTANI[2], NUWAYYIR MOHAMMAD ALSHAMMARI[3], HAZEM ABDULLAH AL-ASMARI[1], ABDULRAHAMAN SALEM ALZAHRANI[2], HUSSEIN HASSAN ALKHAMIS[1], ABDULLAH MOHAMMED ALOWAIFEER[2], & ABDULWAHED FAHAD ALREFAEI[3]*

*[1]Green Sustainability Company for Environmental Services (GSCES), Riyadh, Saudi Arabia *(agrochedad@yahoo.fr)*
[2]The King Abdulaziz Royal Reserve (KARR), Riyadh, Saudi Arabia
*[3]Department of Zoology, College of Science, King Saud University, Riyadh, Saudi Arabia *(afrefaei@ksu.edu.sa)*

Received 23 April 2026, Accepted 13 June 2026

ABSTRACT

Chedad, A., Al-Qahtani, A. M., Alshammari, N. M., Al-Asmari, H. A., Alzahrani, A. S., Alkhamis, H. H., Alowaifeer, A. M., & Alrefaei, A. F. (2026). From coast to desert: First inland record of the Sooty Gull *Ichthyaetus hemprichii* in Saudi Arabia. *Marine Ornithology*, *54*(2), 475–478. http://doi.org/10.5038/2074-1235.54.2.1713

On 22 March 2026, a Sooty Gull *Ichthyaetus hemprichii*, a species normally found in coastal regions of the Red Sea, Arabian Gulf, and western Indian Ocean, was recorded at the Sudair wetlands within the King Abdulaziz Royal Reserve in Saudi Arabia. This site is an inland treated wastewater pond area located at 662 m above sea level. This observation is the first confirmed inland record of the species in Saudi Arabia and the first record within the reserve. The finding extends the known distribution of the species and suggests that occasional inland movements may occur. It highlights the importance of inland protected areas for documenting unusual bird occurrences and improving our understanding of bird distribution in the Arabian region.

Key words: Arabian Peninsula, avifauna, diversity, King Abdulaziz Royal Reserve, migration, wetlands

RÉSUMÉ

Le 22 mars 2026, un Goéland de Hemprich *Ichthyaetus hemprichii,* une espèce généralement observée dans les régions côtières de la mer Rouge, du golfe Arabo-Persique et de l'ouest de l'océan Indien, a été signalé dans les zones humides de Sudair, au sein de la réserve royale du roi Abdulaziz, en Arabie saoudite. Ce site correspond à des bassins intérieurs d'eaux usées traitées situés à 662 m d'altitude. Cette observation constitue la première mention confirmée de l'espèce à l'intérieur des terres en Arabie saoudite, ainsi que la première au sein de cette réserve. Cette découverte élargit l'aire de répartition connue de l'espèce et suggère que des déplacements occasionnels vers l'intérieur des terres peuvent se produire. Elle souligne également l'importance des aires protégées intérieures pour documenter des observations ornithologiques inhabituelles et améliorer la compréhension de la distribution des oiseaux dans la région arabique.

Mots-clés: Péninsule Arabique, avifaune, diversité, réserve royale du roi Abdulaziz, migration, zones humides

OBSERVATIONS

During a bird survey conducted on 22 March 2026 at the Sudair wetlands (wastewater treatment ponds within the industrial area; 25°38′47.90″N, 045°45′56.31″E; 662 m elevation; Fig. 1), a Sooty Gull *Ichthyaetus hemprichii* was observed. The species was confidently identified based on its medium size, dark plumage, sooty-brown coloration, and contrasting white eye crescents (Fig. 2). This species is strongly associated with coastal and nearshore ocean environments, particularly coral islands and sandy shores (Burger et al., 2020; Cramp & Simmons, 1983; del Hoyo et al., 1996; Urban et al., 1986).

In Saudi Arabia, this species is an uncommon resident breeder, with breeding largely restricted to undisturbed offshore islands in the Red Sea, while foraging in adjacent coastal waters. To date, no inland occurrences have been documented (El Ghamdi et al., 2020), making this record from the Sudair wetlands the first documented inland occurrence of the species in Saudi Arabia. Globally, it is distributed along the coasts of the Red Sea, Arabian Sea, and southern Arabian Gulf, as well as the coasts of Pakistan and from eastern South Africa to Kenya. It is regarded as a near-endemic species of the northwestern Indian Ocean, reflecting its strong ecological association with marine and coastal environments (BirdLife International, 2018; Burger et al., 2020).

This gull is considered a partial migrant, with most populations undergoing post-breeding dispersal between September and November, although some populations remain resident year-round in the Red Sea region (del Hoyo et al., 1996; Malling Olsen & Larsson, 2003; Urban et al., 1986). Breeding occurs between April and October, typically in small, loose colonies on coastal or inshore islands, but solitary nesting also occurs occasionally (del Hoyo et al., 1996; Urban et al., 1986). The species is rarely recorded inland or in freshwater environments (Cramp & Simmons, 1983; del Hoyo et al., 1996; Urban et al., 1986). Its diet consists mainly of dead fish and fishery discards, but it may also opportunistically prey on seabird eggs and chicks, as well as turtle hatchlings, making it an important local predator within breeding colonies (del Hoyo et al., 1996; Gallagher et al., 1984; Urban et al., 1986).




Fig. 1. Location of Sooty Gull *Ichthyaetus hemprichii* sightings in Saudi Arabia.

Although the species is currently classified as Least Concern (LC), its global population appears to be decreasing (Boland & Burwell, 2020; Calabrese et al., 2024; Symes et al., 2015). The presence of this individual at Sudair may reflect opportunistic foraging driven by the availability of anthropogenic food resources in artificial habitats such as wastewater treatment ponds. Alternatively, its presence may represent dispersal beyond its typical coastal ranges, underscoring the importance of documenting such atypical



Fig. 2. Sooty Gull *Ichthyaetus hemprichii* at Sudair wetland, Riyadh, King Abdulaziz Royal Reserve, Saudi Arabia, on 22 March 2026: (A) at rest; (B) in flight. Photo credits: Abdelwahab Chedad

occurrences to improve our understanding of this species' ecology and distribution.

The King Abdulaziz Royal Reserve (KARR), which encompasses a wide range of ecosystems and diverse terrain, constitutes a critical area for regional biodiversity, notably avifauna, due to its strategic position along a major migratory flyway between Africa and Asia. Consequently, the reserve may be considered a key bird biodiversity hotspot in Saudi Arabia (Al-Qahtani et al., 2025). Its ecological importance is further supported by several recent noteworthy ornithological records and breeding events within the area, including the occurrence of the Pectoral Sandpiper *Calidris melanotos* (Al-Qahtani et al., 2025); the breeding range expansion of the Black-winged Kite *Elanus caeruleus* (Altalhi et al., 2025); the first evidence of breeding and eastward range expansion of the Mediterranean Short-toed Lark *Alaudala rufescens* (Al-Asmari et al., 2025a); the first confirmed nesting of the Egyptian Nightjar *Caprimulgus aegyptius* (Chedad et al., 2026a); the first record of Rüppell's Vulture *Gyps rueppelli* (Al-Asmari et al., 2025b); the first records of nine vagrant bird species, including raptors, landbirds, and waterbirds (Chedad et al., 2026b); and the first confirmed record of the Red Phalarope *Phalaropus fulicarius* (Chedad et al., 2026c).

ACKNOWLEDGEMENTS

The authors extend their appreciation to the King Abdulaziz Royal Reserve Development Authority and the Green Sustainability Company for Environmental Services (GSCES) in Saudi Arabia for their support and assistance with sample collection. We appreciate the reviewers' comments, which helped us improve our manuscript.

CONFLICT OF INTEREST

The authors declare that no competing interests exist.

FUNDING

This project was funded by King Abdulaziz Royal Reserve Development Authority and conducted by the Green Sustainability Company for Environmental Services, Saudi Arabia.

AUTHOR CONTRIBUTIONS

Observations: AC, AMA, HAA, NMA. Writing—original draft: AC, AFA. Writing—review & editing: AC, AFA, ASA, HHA. All authors read and approved the final version of the manuscript. Resources were provided by ASA, HHA, and AMA.

REFERENCES

Al-Asmari, H. A., Chedad, A., Al-Qahtani, A. M., Alshammari, N. M., Altalhi, M. H., Alzahrani, A. S., Alkhamis, H. H., Alowaifeer, A. M., & Alrefaei, A. F. (2025a). First evidence of breeding and eastward range expansion of the Mediterranean short-toed lark (*Alaudala rufescens*) in King Abdulaziz Royal Reserve, Saudi Arabia. *Biodiversitas*, *26*(8), 4048–4056. https://doi.org/10.13057/biodiv/d260834

Al-Asmari, H. A., Chedad, A., Riad, S., Al-Qahtani, A. M., Alshammari, N. M., Altalhi, M. H., Alzahrani, A. S., Alkhamis, H. H., Alowaifeer, A. M., & Alrefaei, A. F. (2025b). First record of Rüppell's Vulture (*Gyps rueppelli*) in the King Abdulaziz Royal Reserve, Eastern Saudi Arabia. *Vulture News*, *89*(1), 50–56. https://doi.org/10.4314/vulnew.v89i1.7

Al-Qahtani, A. M., Chedad, A., Alshammari, N. M., Altalhi, M. H., Al-Asmari, H. A., Alzahrani, A. S., Alkhamis, H. H., Alowaifeer, A. M., & Alrefaei, A. F. (2025). First confirmed record of *Calidris melanotos* (Vieillot, 1819) Pectoral Sandpiper (Charadriiformes, Scolopacidae), at King Abdulaziz Royal Reserve, Saudi Arabia. *Check List*, *21*(4), 750–754. https://doi.org/10.15560/21.4.750

Altalhi, M. H., Chedad, A., Alshammari, N. M., Al-Qahtani, A. M., Al-Asmari, H. A., Alzahrani A. S., Alkhamis, H. H., Alowaifeer, A. M., & Alrefaei, A. F. (2025). Breeding range expansion of the Black-winged Kite (*Elanus caeruleus*): Evidence from the King Abdulaziz Royal Reserve, Saudi Arabia. *Zoology and Ecology*, *35*(2), 177–183. https://doi.org/10.35513/21658005.2025.2.9

BirdLife International. (2018). *Larus hemprichii. The IUCN Red List of Threatened Species: e.T22694303A132539775*. Retrieved March 25, 2026, from https://dx.doi.org/10.2305/IUCN.UK.2018-2.RLTS.T22694303A132539775.en

Boland, C. R. J., & Burwell, B. O. (2020). Ranking and mapping the high conservation priority bird species of Saudi Arabia. *Avian Conservation & Ecolology*, *15*(2), Article 18. https://doi.org/10.5751/ACE-01705-150218

Burger, J., Gochfeld, M., Kirwan, G. M., & Garcia, E. (2020). Sooty Gull (*Ichthyaetus hemprichii*). In J. del Hoyo, A. Elliott, J. Sargatal, D. A. Christie & E. de Juana (Eds.). *Birds of the world*. Cornell Lab of Ornithology. https://doi.org/10.2173/bow.soogul2.01

Calabrese, L., Riordan, J. A., Lloyd, I. A., Foster, A. D., Collier, T. E., Chambon, J. A., Aljohani, Y. W., Alhamdi, E. A., Beaumont, P. R., Williams, I. D., & Al-Attas, O. (2024). A sea of birds: First bird population assessments in the Saudi Arabian Red Sea. *Frontiers in Marine Science*, *11*, Article 1379601. https://doi.org/10.3389/fmars.2024.1379601

Chedad, A., Alshammari, N. M., Al-Qahtani, A. M., Alhaftaa, A. R., Alzahrani, A. S., Alkhamis, H. H., Alowaifeer, A. M., & Alrefaei, A. F. (2026a). First confirmed nesting record of Egyptian Nightjar *Caprimulgus aegyptius* in Saudi Arabia: King Abdulaziz Royal Reserve. *Applied Ecology and Environmental Research*, *24*(2), 3469–3474. https://doi.org/10.15666/aeer/2402_34693474

Chedad, A., Alshammari N. M., Al-Qahtani, A. M., Riad, S. A., Al-Asmari, H. A., Shahin, A., Althobaiti, A., Alzahrani, A. S., Alkhamis, H. H., Alowaifeer, A. M., & Alrefaei A. F. (2026b). First records of nine vagrant bird species (Aves) from the King Abdulaziz Royal Reserve, Saudi Arabia. *Check List*, *22*(3), 535–548. https://doi.org/10.15560/22.3.535

Chedad, A., Al-Qahtani, A. M., Al-Asmari, H. A., Alzahrani, A. S., Alshammari, N. M., Alkhamis, H. H., Alowaifeer, A. M., & Alrefaei, A. F. (2026c). First confirmed record of the Red Phalarope, *Phalaropus fulicarius* (Linnaeus, 1758) (Charadriiformes, Scolopacidae), at King Abdulaziz Royal Reserve (KARR), Saudi Arabia. *Check List*, *22*(3), 549–554. https://doi.org/10.15560/22.3.549

Cramp, S., & Simmons, K. E. L. (Eds.). (1983). *Handbook of the birds of Europe, the Middle East and North Africa: The birds of the Western Palearctic (Vol. 3, Waders to gulls).* Oxford University Press.

del Hoyo, J., Elliott, A., & Sargatal, J. (Eds.). (1996). *Handbook of the birds of the world (Vol. 3, Hoatzin to auks).* Lynx Edicions.

El Ghamdi, S., Bouzaid, A., & Adarbeh, M. (2020). *Birds of Saudi Arabia I.* Saudi Aramco.

Gallagher, M. D., Scott, D. A., Ormond, R. F. G., Connor, R. J., & Jennings, M. C. (1984). The distribution and conservation of seabirds breeding on the coasts and islands of Iran and Arabia. In J. P. Croxall, P. G. H. Evans, & R. W. Schreiber. (Ed.), *Status and conservation of the world's seabirds*. International Council for Bird Preservation.

Malling Olsen, K., & Larsson, H. (2003). *Gulls of Europe, Asia, and North America*. Christopher Helm.

Symes, A., Taylor, J., Mallon, D., Porter, R., Simms, C., & Budd, K. (2015). *The conservation status and distribution of the breeding birds of the Arabian Peninsula: Regional assessment*. IUCN.

Urban, E. K., Fry, C. H., & Keith, S. (1986). *The birds of Africa* (Vol. 2). Academic Press.